

The Ability Of Klebsiella Pneumonia And Klebsiella Oxytoca To Degrade Oil Waste

by Muna Ali Hussein

Submission date: 24-Jun-2024 11:18AM (UTC+0700)

Submission ID: 2407640682

File name: Natural_-_VOL._2_No._3_Agustus_2024_hal_01-07.pdf (523.8K)

Word count: 2342

Character count: 13980

The Ability Of *Klebsiella Pneumonia* And *Klebsiella Oxytoca* To Degrade Oil Waste

Muna Ali Hussein¹, Yousif Nazzal Hosee², Ayad Ismael Kadhimi³, Umer Jameel Ibrahim⁴

^{1,3}Ministry of Education department, Directorate of Salah Al-Din Education, Tikrit, Iraq.

^{2,4} Environmental Engineering, College of Engineering, Tikrit University, Iraq

corresponding auth: ¹daiemuna20@gmail.com; ²yousif123asd11@gmail.com;

³ayadalkhazrgee1989@gmail.com; ⁴omar.j.ibrahim@tu.edu.iq

8

Article History:

Received: June 20, 2024

Accepted: July 20, 2024

Published: August 31, 2024

Keywords: Oil waste, Biodegradation, wastewater, *Klebsiella pneumonia*, *Klebsiella oxytoca*.

3

Abstract . Crude oil contamination is one of the major environmental problems, it generated processing water pollution by hydrocarbon. Microorganisms have been used to remove or reduce the effects of environmental pollutants as bioremediation agents and the fact it is environmentally friendly. This study isolated two *Klebsiella* strains, *Klebsiella pneumonia* and *Klebsiella oxytoca* from wastewater of the North refineries company in Baiji according to the morphological and biochemical characteristics. This work aimed to evaluate the capacity of two *Klebsiella* strains to degrade the petroleum hydrocarbons efficacy, which excreted from wastewater of the North refineries company. The results showed that the *Klebsiella oxytoca* exhibited a higher hydrocarbon degradation capacity of 67.75% compared to *Klebsiella pneumonia*, which recorded 45.81%. While mixed culture (*Klebsiella oxytoca* and *Klebsiella pneumonia*) was recorded a biodegradation rate of 63.35%. The results showed that efficacy of *Klebsiella oxytoca* as biodegradation for oil wastewater was 30% higher than other strain.

1. INTRODUCTION

Crude oil pollution is a rising environmental concern on a global scale (Singh *et al.*, 2020). Hydrocarbon pollution is a significant environmental issue resulting from several human activities associated with the petrochemical industry (Almansoori *et al.*, 2019). Petroleum wastes as well as crude oil are discharged toward the environment due to activities such as extraction, processing, transportation, storage, accidents, illegal dumping of crude oil wastes, and leaks from deteriorated tanks and pipelines. The principal source of water and soil pollution is the discharge of hydrocarbon pollutants into the environment, whether it occurs accidentally or when untreated before being dispensed. (Adebusoye *et al.*, 2007).

Bioremediation techniques must be developed to degrade hydrocarbons and used to reduce the harmful impacts that petroleum hydrocarbons produce (Singh and Chandra, 2014). The technology of bioremediation is more effective, economical, labor-efficient, reliable, eco-friendly and adaptable. This technology, which works with live things like plants and microbes, is depended on harmful or degradation of toxic organic compounds into harmless compounds (Varjani, 2017; Alfaify *et al.*, 2020)

Muna Ali Hussein daiemuna20@gmail.com

There are two basic methods used in bioremediation schemes: bio-stimulation which involves supplemental nutrients or bio-surfactants and bio-augmentation by adding microbes (Mrozik and Piotrowska-Seget , 2010). It is well known that micro-organisms possessing enzymatic capabilities use petroleum and petroleum hydrocarbons serve as energy and carbon sources (Thapa *et al.*, 2012). Several bacterial species break down hydrocarbons in solution by emulsifying them. They do this by creating biosurfactants, which are active surface agents that help the cells stick to the substrate (Hassanshahiant *et al.*, 2012). Bhattacharya *et al.* ,2003 reported on the catabolic ability of *Klebsiella* strains are gram-negative bacteria to break down hydrocarbons, especially polyaromatic hydrocarbon (PAH). The purpose of this work is to isolate bacterial *Klebsiella* strains that can be used in bioremediation of hydrocarbon contaminated water, and describe the ability of two *Klebsiella* strains and efficiently to degrade crude oil and its related products under laboratory conditions. There may be potential uses for this research in the remediation of hydrocarbon contamination.

2. METHODOLOGY

2.1. Sample collection

Samples of industrial wastewater polluted with oil waste were collected from various places in the water flow from which oil waste is released to the North Refineries Company in Al-Baiji, transported to the laboratory in sterilized glass bottles and stored at 4 C°, until they were used to isolate the bacteria.

2.2. Isolation and identification of bacteria

Bacteria were isolated from oil wastewater by using dilution method, which was prepared from 1 ml of the wastewater sample in a test tube with 9 ml of normal saline solution to obtain the required concentration, then it was taken from a dilution of 103 and spread on nutrient agar and incubate at 37 C° for an hour, according to (Olukunle ,2013). Bacteria were identified according to the morphological characteristics, the shape and cell size, and their Gram stain interaction test when compared to Bergey's manual (Murray *et al.*, 2003), and using biochemical tests (Catalase, oxidase enzyme production, acylation of gelatin, H₂S gas production, starch decomposition, nitrate consumption, motility test, indole test, hemolysis, ability to ferment sugars). Two species of bacteria belonging to the *Klebsiella* genus were identified and the isolated were preserved on nutrient broth at refrigerator temperature.

2.3 Measuring of quantitative loss% of hydrocarbon compounds

The measure of oil waste decomposition rate by using the gravimetric method to calculate the percentage of hydrocarbon consumption by calculating the Residue amount of hydrocarbon

compounds by measuring the difference between the amount of oil waste weight added to the culture medium used before and after the bacterial growth used in the study. The percentage rate of decomposition by Separately and synergistically was calculated for the bacterial isolates, and statistical analyses were performed for these results.

The infrared spectra of bacterial samples analyzed belonging to the *Klebsiella* genus in this study were measured by Shimadzu FT-IR 8400 spectrophotometer ranged 400-4000cm⁻¹ using KBr carried out at Chemistry department / College of Science/Tikrit University.

3. RESULTS AND DISCUSSION

Two *Klebsiella* species were used to analyze petroleum residues in the waste water, they were distinguished from each other using the tests outlined in Table 1. The findings indicated the presence of *Klebsiella* species. The isolates exhibited characteristics of Gram-negative, facultative anaerobic, non-motile short rods, non-sporing. Biochemical tests revealed that the isolates were catalase positive, oxidase negative, methyl red test negative, Vogus-Proskauer and glucose test positive, and Urease test positive. Additionally, the isolates displayed diverse reactions to other biochemical tests. The two medically significant species, *Klebsiella pneumoniae* and *Klebsiella oxytoca*, can be distinguished based on their ability to synthesize indole from tryptophan. *K. oxytoca* is classified as indole positive. (Maslow *et al.*, 2003).

Table 1. Biochemical tests of the *Klebsiella* species at the present study.

Bacteria isolation	Biochemical tests											
	Gram stain	Catalase	Oxidase	H ₂ S	Gelatin	Nitrate	Urease	Methyl red	Indole test	Motion test	Glucose	Lactose test
<i>Klebsiella pneumoniae</i>	-	+	-	-	-	+	+	-	-	-	+	+
<i>Klebsiella oxytoca</i>	-	+	-	+	+	+	+	-	+	-	+	+

The removal of hydrocarbons through microbial degradation has been regarded as an efficient and eco-friendly method, a various microorganisms isolated from contaminated locations can degrade hydrocarbons under aerobic and facultative anaerobic conditions (Kumar & Ghosh, 2022). The *Klebsiella* bacteria used were studied for their ability to consume

hydrocarbon content. It was found that the *Klebsiella oxytoca* exhibited a higher hydrocarbon degradation capacity compared to *Klebsiella pneumonia* and a mixed culture (*Klebsiella oxytoca* + *Klebsiella pneumonia*), which recorded a lower degradation rate. The rate ability of the *Klebsiella oxytoca* to the hydrocarbon degradation is 67.75% whereas *Klebsiella pneumonia* is 45.81%, while the mixed culture of bacteria was recorded of 63.35% for the hydrocarbon degradation (Table 2).

Table 2. Hydrocarbon degradation percentage in waste water using *Klebsiella*.

Bacteria isolates	Wt. of untreated Crude oil (cm ³)	Remaining wt. of un-degradation (cm ³)	Wt. of degradation petroleum waste (cm ³)	Degradation rate %
<i>Klebsiella pneumonia</i>	0.8447	0.4577	0.3870	45.81
<i>Klebsiella oxytoca</i>	0.8447	0.2724	0.5723	67.75
<i>Klebsiella pneumonia</i> + <i>Klebsiella oxytoca</i>	0.8447	0.5352	0.6335	63.35

Chamkha et al. (2011) observed that *Klebsiella oxytoca* has the ability to break down a diverse spectrum of aliphatic hydrocarbons, specifically those with carbon chain lengths ranging from C13 to C30. In their study, Mohamed et al. (2012) demonstrated that *K. oxytoca*, obtained from crude petroleum oil samples, exhibited the ability to metabolize four distinct polycyclic aromatic hydrocarbons (PAHs): benzene, phenanthrene, pyrene, and fluoranthene. The infrared spectroscopy technique (IR) was used to determine the various bonds in the compounds. It was found that the chemical bonds, including stretching vibrations of C-H and C-C bonds with less frequency appearance, fall within the range of 4000-1300 cm⁻¹. The second effect is bending vibrations, which involve changes in the positions of bonds without their disappearance. All the oil residue samples gave similar absorption spectra, indicating that the groups and bonds in the compounds are similar.

Several studies have confirmed the use of this technique in studying crude oil degradation (Al-Asadi, 2000; Al-Jubouri, 2005). Additionally, (Al-Obaidi, 2003) pointed out changes in the positions of atoms and bonds, indicating the role of bacterial isolates in the degradation of aromatic compounds. Multiple peaks were observed for aromatic hydrocarbons and phenols in crude oil treated with *Klebsiella pneumonia* and *Klebsiella oxytoca*, as shown in Figure1,

compared to Figure (2,3) which represent the degradation of oil residue samples by the bacterial species *Klebsiella pneumonia* and *Klebsiella oxytoca*, and synergistically as mixed culture using this technique. The results show a significant change in the absorption positions of the bonds, but differences in the intensity of the peaks in the overall shape of the spectra were observed. The two spectra mentioned above were sufficient to demonstrate that there were differences in the spectra for degradation using this biological method.




Figure 1. IR spectrum of the oil residue sample before the treatment by bacterial isolates.




Figure 2. IR spectrum of the oil residue sample after treatment by *Klebsiella oxytoca*.




Figure 3. IR spectrum of the oil residue sample after treatment by *Klebsiella pneumonia*.

4. CONCLUSIONS

The degradation efficacy of *Klebsiella oxytoca* for oil wastewater like petroleum hydrocarbons was a 30% greater than the *Klebsiella pneumonia*. While a degradation rate of 63.35% was recorded for the hydrocarbon content degradation in the mixed culture of bacteria. This result showed that these *Klebsiella* strains would need to break down petroleum hydrocarbon contaminants as the eco-friendly technology clean-up (bioremediation). So, strains producing bio-surfactants may play an important role in petroleum hydrocarbon biodegradation, it enhancing biodegradation process.

5. ACKNOWLEDGEMENTS

The authors thank the North Refineries Company in Al-Baiji and Chemistry department at the College of Science / Tikrit University for their support of this work.

REFERENCES

- Adebusoye, S. A., Ilori, M. O., Amund, O. O., Teniola, O. D., & Olatope, S. O. (2007). Microbial degradation of petroleum hydrocarbons in a polluted tropical stream. *World Journal of Microbiology and Biotechnology*, 23, 1149-1159. <https://doi.org/10.1007/s11274-007-9345-3>
- Al-Asadi, M. J. (2000). Synthesis and evaluation of new demulsifiers for use in the fields of Southern Petroleum Company. (Master's thesis). College of Science, Basrah University, Iraq.
- Alfaify, A. M., Mir, M. A., & Alrumman, S. A. (2022). *Klebsiella oxytoca*: An efficient pyrene-degrading bacterial strain isolated from petroleum-contaminated soil. *Archives of Microbiology*, 204(5), 248. <https://doi.org/10.1007/s00203-022-02850-9>
- Al-Jubouri, Y. H. A. (2005). The biodegradation of some types of crude oil by the action of the germs isolated from soils in North Iraqi refineries. (Doctoral dissertation). College of Education for Pure Science, Tikrit University, Iraq.
- Almansoori, A. F., Talal, A., Al-Yousif, N. A., & Hazaimah, M. (2019). Isolation and identification of microbial species for hydrocarbon degradation in contaminated soil and water. *Plant Archives*, 19(1), 971-977.
- Al-Obaidi, I. M. (2003). Study of biological degradation and infrared spectrum of samples of Kirkuk crude oil treated with nitrogen-fixing cyanobacteria. *Tikrit Journal of Pure Sciences*, 1, 52-66.
- Bhattacharyya, J. K., & Shekdar, A. V. (2003). Treatment and disposal of refinery sludges: Indian scenario. *Waste Management & Research*, 21(3), 249-261. <https://doi.org/10.1177/0734242X0302100309>

- Hassanshahian, M., Emtiazi, G., & Cappello, S. (2012). Isolation and characterization of crude-oil-degrading bacteria from the Persian Gulf and the Caspian Sea. *Marine Pollution Bulletin*, 64(1), 7-12. <https://doi.org/10.1016/j.marpolbul.2011.11.006>
- Kumar, S., & Ghosh, A. (2022). Fractional-order identification and synthesis of equivalent circuit for electrochemical system based on pulse voltammetry. In *Fractional-Order Design* (pp. 373-402). Academic Press.
- Maslow, J. N., Brecher, S. M., Adams, K. S., Durbin, A., Loring, S., & Arbeit, R. D. (1993). Relationship between indole production and differentiation of *Klebsiella* species: Indole-positive and -negative isolates of *Klebsiella* determined to be clonal. *Journal of Clinical Microbiology*, 31(8), 2000-2003.
- Mrozik, A., & Piotrowska-Seget, Z. (2010). Bioaugmentation as a strategy for cleaning up soils contaminated with aromatic compounds. *Microbiological Research*, 165(5), 363-375. <https://doi.org/10.1016/j.micres.2009.08.001>
- Olukunle, O. F. (2013). Characterization of indigenous microorganisms associated with crude oil-polluted soils and water using traditional techniques. *Microbiology Journal*, 3(1), 1-11.
- Singh, H., Bhardwaj, N., Arya, S. K., & Khatri, M. (2020). Environmental impacts of oil spills and their remediation by magnetic nanomaterials. *Environmental Nanotechnology, Monitoring & Management*, 14, 100305. <https://doi.org/10.1016/j.enmm.2020.100305>
- Singh, K., & Chandra, S. (2014). Treatment of petroleum hydrocarbon polluted environment through bioremediation: A review. *Pakistan Journal of Biological Sciences*, 17(1), 1-8. <https://doi.org/10.3923/pjbs.2014.1.8>
- Thapa, B., Kumar, K. C. A., & Ghimire, A. (2012). A review on bioremediation of petroleum hydrocarbon contaminants in soil. *Kathmandu University Journal of Science, Engineering and Technology*, 8(1), 164-170.
- Varjani, S. J. (2017). Microbial degradation of petroleum hydrocarbons. *Bioresource Technology*, 223, 277-286. <https://doi.org/10.1016/j.biortech.2016.10.037>
- Younis, B. M., & Saeed, I. O. (2023). Concentration of heavy metals in soil contaminated with crude oil at two Iraqi sites according to environmental indices of pollution. *Nativa*, 11(4), 558-565. <https://doi.org/10.31413/nat.v11i4.16521>

The Ability Of Klebsiella Pneumonia And Klebsiella Oxytoca To Degrade Oil Waste

ORIGINALITY REPORT

9%

SIMILARITY INDEX

6%

INTERNET SOURCES

4%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | S. Bilen Ozyurek, I. Seyis Bilkay.
"Biodegradation of petroleum by Klebsiella pneumoniae isolated from drilling fluid",
International Journal of Environmental
Science and Technology, 2017
Publication | 2% |
| 2 | www.mdpi.com
Internet Source | 1% |
| 3 | www.plantarchives.org
Internet Source | 1% |
| 4 | journal.widyakarya.ac.id
Internet Source | 1% |
| 5 | uir.unisa.ac.za
Internet Source | 1% |
| 6 | Submitted to Rancho Santiago Community
College District
Student Paper | 1% |
| 7 | Submitted to Tikrit University
Student Paper | 1% |

8

journal-stiayappimakassar.ac.id

Internet Source

1 %

9

dokumen.pub

Internet Source

<1 %

10

link.springer.com

Internet Source

<1 %

11

Linda U. Obi, Harrison I. Atagana, Rasheed A. Adeleke. "Isolation and characterisation of crude oil sludge degrading bacteria", SpringerPlus, 2016

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On