

Phytochemical Investigation and GC-MS Analysis of Tribulus Terrestris L. Cultivated in Iraq

by Amani Amer Tawfeeq

Submission date: 02-Aug-2024 09:46AM (UTC+0700)

Submission ID: 2426070033

File name: VOL.2_SEPTEMBER_2024_HAL_22-29.docx (673.48K)

Word count: 2373

Character count: 14424



Phytochemical Investigation and GC-MS Analysis of *Tribulus Terrestris* L. Cultivated in Iraq

Amani Amer Tawfeeq^{1*}, Tahany Amir Tawfeeq², Zahraa A.E Al Naqqash³, Zlinah Alaa Kamal⁴

^{1,2,3,4} Department of Pharmacognosy and Medicinal Plants, College of Pharmacy, Mustansiriyyah University, Baghdad, Iraq
dr.amanitawfeeq@uomustansiriyah.edu.iq^{1*}, tahaniamir@uomustansiriyah.edu.iq²,
zahraa_abd_alelah@uomustansiriyah.edu.iq³, zinah85.za@gmail.com⁴

Author correspondence: dr.amanitawfeeq@uomustansiriyah.edu.iq

Abstract: *Tribulus terrestris* L. Gokshur in Sanskrit, is an annual herbaceous, little prostrate, silky from the Zygophyllaceae family. It is found in extensive, arid, and scorching regions of India. is used in traditional folk medicines as an aphrodisiac, astringent, lithotriptic, diuretic, and urinary disinfectant. However, uneven, pinnae, elliptical leaves. Sprinkled, round, compressed, five-cornered, transverse partitioned carpel fruits with many seeds ripen in July. Pieces of the cylindrical-fibrous root are 7-18 cm and 0.3-0.7 cm wide. According to reported studies, the investigations indicated the presence of the saponins :gitogenin, tigogenin, neogitogenin, hecogenin, and flavonoid tribuloside was kaempferol-3-glucoside. Furthermore, *Tribulus terrestris* plays a vital role in hepatotoxicity and is greatly reduced by *Tribulus* extracts. Similarly, *T. terrestris* plays a remarkable role in reducing the elevated level of serum glucose, triglycerides, VLDL, and cholesterol. This study aims to conduct a preliminary analysis of the phytochemicals of *Tribulus* leaves cultivated in Iraq and to determine the active constituent of the ethanolic extract of the leaves using Gc-Mass. Qualitative detection of herbal leaves in Iraqi soil, investigated by GC-MASS revealed important constituents. The funding shows a new compound; dihydrocoumarin and other major constituents neophytadiene and alpha-tocopherol.

Keywords: *Tribulus*, Mass, dihydrocoumarin, alpha-tocopherol, phytochemical.

1. INTRODUCTION

Tribulus terrestris, known as Gokshur (Sanskrit); is distributed along wide dry, and hot geographic regions in India and It is grown in worm regions in southern Eurasia and Africa. The morphological characteristics of the shrub are characterized by a small prostrate - silky hairy shrub, with an aerial part height of 10-60 cm(1). Leaves are opposite, may be unequal, pinnae, and elliptical. Contrasting, uneven, pinnae, elliptical leaves. Sprinkled, round, compressed, five-cornered, transverse partitioned carpel fruits with many seeds ripen in July. Pieces of the cylindrical-fibrous root are 7-18 cm in length and 0.3-0.7 cm wide(2). Occasionally, investigations were reported that *T. terrestris* has different bioactivity secondary metabolites of saponins, alkaloids, flavonoids, and tannins. The reported studies revealed that the presence of the saponins of gitogenin, tigogenin, neogitogenin, and hecogenin, the quantity of main flavonoids is approximately 1.5 times that of main saponins(3)(4). The aerial parts of *T. terrestris* also possess flavonoids; kaempferol and rutin, and its glycosides; tribuloside, kaempferol-3-glucoside, and rutinoid(5)(6). Chemical constituents harmane and norharmane presented in *T. terrestris* as alkaloids^[11]. Tian Shung et al. isolated two new compounds,

tribulusterine, terrestribisamide, and, known compounds, β -sitosterol, ferulic acid, and vanillin from the dried fruits(7). *T. terrestris* in folk or unofficial medicines, is used mainly as a diuretic, aphrodisiac, and astringent(8). However, its pharmacological activity reveals a significant effect of essential oils and elements, the high concentration of potassium salts, on its diuretic activity(9). The *Tribulus* extracts showed incredible effects against acetaminophen-induced hepatotoxicity(10) and Improvement in fertility(11). Furthermore, the herb also exhibited a pro-erectile effect after oral administration on rabbit corpus cavernous smooth muscle(12). Likewise, *T. terrestris* has a remarkable role in reducing the high level of serum glucose and decreasing serum triglyceride, very low-density lipoprotein (VLDL), and serum cholesterol(13). The aim of this study involves the Phytochemical investigation of *Tribulus Terrestris* leaves grown in Iraq and the analysis of the active constituent of the leaves ethanolic extract using Gc-Mass.

2. MATERIALS AND METHODS:

Plant material and chemical materials:

The plant material was prepared from the Leaves of *Tribulus terrestris* as shown in Figure 1. They were collected from Al Zahwraa Park Iraq\ Baghdad and authenticated by Dr. Ibrahim Salih at the College of Pharmacy-Mustansiriyah University. The leaves were collected during October/ 2023 and dried under shade conditions at room temperature for 14 days before grinding commercial electric blender and weighing as a powder.



Figure 1: *Tribulus terrestris* cultivated in Iraq

1. Equipment and Chemicals:

Ethanol (95-97%) was purchased from Merck (Darmstadt, Germany), and A rotary evaporator (BÜCHI Rotavapor R-205, Swiss) was used for the dryness of the extracts. Apparatus ,GC/MS -SHIMADZU -QP-2010 ULTRA-Scan mode -ACQ .

2. Phytochemical investigations for *Tribulus terrestris* leaves:

The preliminary chemical tests to screen bioactive compounds and GC -MS analysis for ethanolic extract were performed as follows

Preliminary phytochemical Tests of *Tribulus terrestris* :

Qualitative analysis of leaves was performed according to standard procedures to indicate preliminary analysis of chemical constituents(14), (15),(16).

Test for screening of Saponins (Foam Test):

Approximately one mL of Methanolic extract was mixed with (5mL) of distilled water in a test tube and then shaken vigorously, the persistent foam at least 1cm in height indicates the presence of saponin.

1. Test for screening of Terpenoids (Salkowski Test):

About 4mL of alcoholic extract was mixed with chloroform (2mL), then carefully added 3mL of concentrated sulphuric acid. A reddish–brown color is formed to indicate the presence of terpenoids.

2. Benedict's test:

Benedict's reagent (5mL) was added in the test tube to the methanolic extract (1mL) and then the mixture was heated in a boiling water bath for 10 minutes. The green precipitate appearance is an indication of the presence of the reducing sugar.

3. Test for screening of Phenols:

Alcoholic extract (0.5 mL) was treated with 3 to 4 drops of ferric chloride. The appearance of an intense green color indicates the presence of phenols.

4. Wagner's reagent for screening of Alkaloids:

Approximately 3 mL of Alcoholic extract acidify with a few drops of hydrochloric acid. Then a few drops of Wagner's reagent added and the resulting color of the precipitate was reported.

5. Tests for screening of Flavonoids:

Approximately 3mL of alcoholic KOH was added to 1 mL of methanolic leaf extract. A yellow color indicates a relevant result.

6. Test for screening of Tannins:

About 0.2g of sample was dissolved in (10ml) of distilled water and filtered. Then a Solution of (1% FeCl₃) was added. the appearance of dark green, dark blue, or black color related to the presence of tannins.

7. Test for screening of Coumarins:

A few drops of methanol extract from leaves of *Tribulus terrestris* were placed on filter paper, adding two drops of NaOH (1N) solution, and then the filter paper was detected under UV light at 366nm for the presence of yellow to blue fluorescence.

Extraction of leaves *Tribulus terrestris*:

For the obtaining of extracts, the obtained powdered leaves of *T.terrestris* (50 g) were extracted with ethanol 350 mL(16). Extraction was carried out using a Soxhlet extractor, and the collected extracts were then filtered using a filter paper and concentrated using a rotary evaporator, The obtained extract was stored at 4 °C in dark containers for analysis with Gas Chromatography /Mass.

GC-MS analysis:

Gas Chromatography /Mass was performed and applied to detect phytochemicals in the ethanol extract. The GC-MS conditions were: the carrier gas was helium, the injection volume was 1µL, and the split ratio was 2.0, the Injection temperature was 250°C, Column temperature was about 80 °C to 310 °C at a rate of 10C° /min.

3. RESULTS AND DISCUSSIONS

Phytochemicals investigation:

The preliminary tests are essential in predicting different phytochemicals in the plant extracts.

Table 1. Phytochemical screening of Iraqi *Tribulus terrestris*

Chemical Tests	Positive Indicators	Results
Saponin Test	Foam observed 1cm in high	+ ve
Tannins Test	Dark green, dark blue, or black color	+ ve
Flavonoid Test	Yellow color	+ ve
Alkaloid Test	Reddish-brown precipitate	_ ve

**PHYTOCHEMICAL INVESTIGATION AND GC-MS ANALYSIS OF *TRIBULUS TERRESTRIS*
L. CULTIVATED IN IRAQ**

Terpenes Test	Reddish-brown color	+ ve
Coumarin test	yellow to blue fluorescence	+ ve
Benedicts test	green precipitate	+
		ve

The results (table 1) showed the presence of different phytochemicals such as flavonoids, tannins, terpenes, coumarins, saponins, and reducing sugar with the absence of alkaloids that are vital in the biological activities of *T.terrestris* plant. These findings were agreed with reported literature in which positive results for flavonoids, carbohydrates, and terpenes tests were obtained (17).

GC/MS Analysis

Analysis by GC-MASS for the qualitative detection of important constituents and determination of bioactive chemical components identified in leaves ethanol extract shows different components and groups of terpenes and steroids as shown in (Figure 2) and (Table 2) shows the major components identified in the ethanolic extract of *T.terrestris* leaves. The GC-MS chromatogram shows the presence of more different peaks representing chemical constituents. The presence of dihydrocoumarin at a retention time of 17.079 min, neophytadiene (R.time 23.726 min) , alpha-tocopherol at (R.time 24.463), and Silicic acid (R.time 43.28min). The chromatogram can be seen in (Figure 2) with peaks retention time and Similarity index (SI).




Figure 2: GC/MS chromatogram of methanolic extract of *T.terrestris* leaves

Table 2: GC -MS analysis of main chemical constituents of *T.terrestris* leaves.

Peak NO.	Name	R. time	SI	Area %
1	Di hydrocoumarin	17.079	95	59.07
2	2,4-Di-tert-butylphenol	17.754	97	0.73
3	Neophytadiene	23.726	96	1.71
4	alpha-Tocopherol	24.463	97	0.74
5	4,4'-Ethylenebis(2,6-di-tert-butylphenol)	36.936	93	0.57
6	Silicic acid, diethyl bis(trimethylsilyl) ester	43.282	96	0.77
7	Phenol, 2,4-bis(1,1-dimethylethyl)-, phosphite	45.129	95	23.06
8	Silane, dimethyl(2-naphthoxy)heptyloxy-	49.041	97	13.35

Dihydrocoumarins, one of the coumarin derivatives, have been reported to influence antioxidant and anti-inflammatory properties. Additionally, they function as an HDAC inhibitor and increase DNA damage sensitivity by inhibiting Rad52(18). Neophytadiene is a diterpene; 7,11,15-trimethyl-3methylidenehexadec-1-ene. Its role an antimicrobial agent, and anti-inflammatory agent (19). Alpha-tocopherol also Enhances the immune system and prevents the formation of blood clots, it aids in the prevention of cellular damage caused by free radicals (antioxidant effect). Vitamin E is currently under investigation in protecting and preventing cancer(20). Silicic acid (H_4SiO_4), an accessible silicon for humans and animals has been disregarded thus far. Silicon is associated with effects such as bone mineralization, collagen synthesis, skin, hair, and nail health, atherosclerosis, Alzheimer's disease, immune system enhancement, and certain other disorders or pharmacological effects have all been linked to silicon(21).

Conclusions

This study proved that the Iraqi *T.terrestris* contains various secondary metabolites coumarins, flavonoids, terpenes, carbohydrates, and saponin. Dihydrocoumarins is a new compound revealed in GC-Mass analysis that is considered avital as anti-inflammatory and anti-oxidants. other compounds Vitamin E (tocopherol), and Neophytadiene revealed the presence of minor compounds in *T.terrestris* leaves. Outlook, this study is required for further investigations and structural characterizations of phytochemicals in all plant parts.

Acknowledgment

We appreciate the logistical assistance by Mustansiriyah University of Pharmacy - Pharmacognosy and Medicinal Plant Department.

REFERENCES

- Adaikan, P. G., Gauthaman, K., Prasad, R. N., & Ng, S. C. (2000). Proerectile pharmacological effects of *Tribulus terrestris* extract on the rabbit corpus cavernosum. *Annals of the Academy of Medicine, Singapore*, 29(1), 22–26.
- Akram, M., Asif, H. M., Akhtar, N., Shah, P. A., Uzair, M., & Shaheen, G., et al. (2011). *Tribulus terrestris* Linn.: A review article. *Journal of Medicinal Plants Research*, 5(16), 3601–3605.
- Al-Ali, M., Wahbi, S., Twaij, H., & Al-Badr, A. (2003). *Tribulus terrestris*: Preliminary study of its diuretic and contractile effects and comparison with *Zea mays*. *Journal of Ethnopharmacology*, 85(2–3), 257–260.
- Chhatre, S., Nesari, T., Somani, G., Kanchan, D., & Sathaye, S. (2014). Phytopharmacological overview of *Tribulus terrestris*. *Pharmacognosy Reviews*, 8(15), 45.
- Chen, C. C., Huang, J. S., Wang, T. H., Kuo, C. H., Wang, C. J., & Wang, S. H., et al. (2017). Dihydrocoumarin, an HDAC inhibitor, increases DNA damage sensitivity by inhibiting Rad52. *International Journal of Molecular Sciences*, 18(12).
- Eldalawy, R., Al-Ani, W. M. K., & Kareem, W. A. (2021). Phenotypic, anatomical and phytochemical investigation of Iraqi *Silybum marianum*. In *Journal of Physics: Conference Series* (p. 22029). IOP Publishing.
- Ibrahim, N. (2015). Phytochemical investigation and antioxidant activity of Iraqi *Tribulus terrestris*. *Baghdad International Journal of Pharmaceutical Sciences*, 24(1). Available from: <https://bijps.uobaghdad.edu.iq/index.php/bijps/article/view/380>
- Jurkić, L. M., Cepanec, I., Pavelić, S. K., & Pavelić, K. (2013). Biological and therapeutic effects of ortho-silicic acid and some ortho-silicic acid-releasing compounds: New perspectives for therapy. *Nutrition & Metabolism*, 10(1), 2.
- Kamenov, Z., Fileva, S., & Kalinov, K. (2015). Evaluation of the efficacy and safety of *Tribulus terrestris* in male sexual dysfunction—a prospective, randomized, double-blinded, placebo-controlled clinical trial. *Maturitas*, 81(1), 208.
- Kavitha, P., Ramesh, R., Bupesh, G., Stalin, A., & Subramanian, P. (2011). Hepatoprotective activity of *Tribulus terrestris* extract against acetaminophen-induced toxicity in a freshwater fish (*Oreochromis mossambicus*). *In Vitro Cellular & Developmental Biology-Animal*, 47, 698–706.
- Kostova, I., & Dinchev, D. (2005). Saponins in *Tribulus terrestris*—chemistry and bioactivity. *Phytochemistry Reviews*, 4, 111–137.

- Li, M., Qu, W., Wang, Y., Wan, H., & Tian, C. (2002). Hypoglycemic effect of saponin from *Tribulus terrestris*. *Zhong Yao Cai = Zhongyaocai = Journal of Chinese Medicinal Materials*, 25(6), 420–422.
- Louveaux, A., Jay, M., El Hadi, O. T. M., & Roux, G. (1998). Variability in flavonoid compounds of four *Tribulus* species: Does it play a role in their identification by desert locust *Schistocerca gregaria*? *Journal of Chemical Ecology*, 24, 1465–1481.
- Rizvi, S., Raza, S. T., Ahmed, F., Ahmad, A., Abbas, S., & Mahdi, F. (2014). The role of vitamin E in human health and some diseases. *Sultan Qaboos University Medical Journal*, 14(2), e157–e165.
- Rzhepakovsky, I. V., Areshidze, D. A., Avanesyan, S. S., Grimm, W. D., Filatova, N. V., & Kalinin, A. V., et al. (2022). Phytochemical characterization, antioxidant activity, and cytotoxicity of methanolic leaf extract of *Chlorophytum comosum* (Green Type) (Thunb.) Jacq. *Molecules*, 27(3).
- Saleh, N. A. M., Ahmed, A. A., & Abdalla, M. F. (1982). Flavonoid glycosides of *Tribulus pentandrus* and *T. terrestris*. *Phytochemistry*, 21(8), 1995–2000.
- Tawfeeq, A. A., Ali, S. H. (2022). Isolation and structural characterization of quercetin 3-O-rhamnoside and essential oil estimation from leaves of Iraqi *Cupressus sempervirens* L (Conference Paper). *Iraqi Journal of Pharmaceutical Sciences (P-ISSN 1683-3597 E-ISSN 2521-3512)*, 31(Suppl.), 121–130.
- Tawfeeq, A. A., Mahdi, M. F., Abaas, I. S., & Alwan, A. H. (2018). Isolation, quantification, and identification of rosmarinic acid, gas chromatography-mass spectrometry analysis of essential oil, cytotoxic effect, and antimicrobial investigation of *Rosmarinus officinalis* leaves. *Asian Journal of Pharmaceutical and Clinical Research*, 11(6), 126–132.
- Tyagi, P., & Ranjan, R. (2023). Comparative study of the pharmacological, phytochemical and biotechnological aspects of *Tribulus terrestris* Linn. and *Pedaliu murex* Linn: An overview. *Acta Ecologica Sinica*, 43(2), 223–233.
- Wu, T. S., Shi, L. S., & Kuo, S. C. (1999). Alkaloids and other constituents from *Tribulus terrestris*. *Phytochemistry*, 50(8), 1411–1415.
- Xu, Y. X., Chen, H. S., Liu, W. Y., Gu, Z. B., & Liang, H. Q. (1998). Two sapogenins from *Tribulus terrestris*. *Phytochemistry*, 49(1), 199–201.

Phytochemical Investigation and GC-MS Analysis of Tribulus Terrestris L. Cultivated in Iraq

ORIGINALITY REPORT

10%

SIMILARITY INDEX

6%

INTERNET SOURCES

6%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|-----|
| 1 | rogersgroup.northwestern.edu
Internet Source | 1 % |
| 2 | Kranti Kamble, Kirti Laddha. "A Method of Analysis of Asafoetida Marketed Samples for its Umbelliferone Content", Research Square Platform LLC, 2022
Publication | 1 % |
| 3 | Xin Fang, Yanan Liu, Jingyi Xiao, Cunqiang Ma, Youyi Huang. "GC-MS and LC-MS/MS metabolomics revealed dynamic changes of volatile and non-volatile compounds during withering process of black tea", Food Chemistry, 2023
Publication | 1 % |
| 4 | elibrary.tucl.edu.np
Internet Source | 1 % |
| 5 | Submitted to Universitas Sebelas Maret
Student Paper | 1 % |
| 6 | Submitted to Universiti Teknologi MARA
Student Paper | 1 % |

7	Rasha Eldalawy, Noor Mohsin Naser, Zahraa A. E. Al Naqqash. "Antimicrobial and antioxidant activity of Iraqi cupressuss empervirens cones", AIP Publishing, 2020 Publication	1 %
8	e-sciencecentral.org Internet Source	1 %
9	eprints.skums.ac.ir Internet Source	1 %
10	Submitted to 2642 Student Paper	1 %
11	Submitted to Guru Jambheshwar University of Science & Technology Student Paper	1 %
12	Submitted to Higher Education Commission Pakistan Student Paper	1 %
13	repository.ush.edu.sd:8080 Internet Source	1 %
14	P. Kavitha, R. Ramesh, G. Bupesh, A. Stalin, P. Subramanian. "Hepatoprotective activity of Tribulus terrestris extract against acetaminophen-induced toxicity in a freshwater fish (Oreochromis mossambicus)", In Vitro Cellular & Developmental Biology - Animal, 2011	1 %

Exclude quotes	On	Exclude matches	< 1%
Exclude bibliography	On		

Phytochemical Investigation and GC-MS Analysis of Tribulus Terrestris L. Cultivated in Iraq

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/0

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8